LETTERS TO THE EDITOR.

[*The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of* NATURE. *No notice is taken of anonymous communications.*]

The Ionisation of Gases in the Schumann Region.

IN NATURE for June 12, Prof. Lyman discusses the evidence relating to the ionisation of air in the Schumann region, and concludes that ionisation of air can be produced by wave-lengths longer than λ1700.

The reasons why I consider that λ1350 is nearer to the limiting wave-length at which the ionisation of air sets in are as follows:—Using a discharge in hydrogen as a source of ultra-violet light and transmitting it through a quartz window (0·3 mm. thick), I was unable to get any ionisation in filtered air. I only obtained big effects with a certain piece of fluorite as the window. Another piece of fluorite which did not transmit the ionising light was transparent to about λ1350. Prof. Lyman's researches show that the hydrogen discharge emits very intense ultra-violet light distributed over a large number of wave-lengths between λ1200 and λ1600. Hence, with thin quartz, there was plenty of light available down to λ1450, but it produced no effect in my experiments. Similarly, wave-lengths down to about λ1350 produced no appreciable effects.

Lenard and Ramsauer used a very intense aluminium spark as their source of light, and found that the light from it transmitted through fluorite produced enormous ionisation in air. On the other hand, the light when passed through quartz did not produce any effect. According to Prof. Lyman's photographs, the wave-lengths available from the Al spark in air are a strong group of lines near λ1300, some weak lines near λ1500, and strong lines near λ1600 and λ1720–λ1800. Thin quartz cuts out the group λ1300, but allows the others to pass. We are not told explicitly whether the spark in Lenard and Ramsauer's researches was ever placed close to the quartz window to avoid air absorption; if so, the λ1500 and λ1600 groups would be effective. Fluorite, on the other hand, transmits the λ1300 group as well, and Prof. Lyman considers that the ionisation observed is probably due to these lines. He points out that my interpretation of his remarks, viz. that λ1300 represents the longest wave-lengths which are effective in ionising air, does not represent his views correctly. He considers that Bloch's recent work on the ionisation of air by a mercury lamp indicates that wave-lengths longer than λ1750 are effective.

I expect it will be agreed that by air we mean the usual mixture of oxygen and nitrogen free from all the more condensable gases. Lenard and Ramsauer found that ordinary dust-free air was certainly ionised by the light transmitted through quartz from their powerful source. It was only when very drastic methods of purification were adopted that the air was no longer ionised by the light transmitted through quartz. Although Bloch used dust-free air in his experiments, there is no evidence that he took the rigorous precautions which Lenard and Ramsauer assert are necessary to get rid of all the impurities which give rise to ionisation with comparatively long wave-lengths. In Bloch's arrangement, the mercury lamp was totally immersed in the stream of air, and consequently all the light emitted was available for ionisation, and hence the traces of impurities have every chance to be ionised. Bloch does not give any details, but I think the supports, insulated wires, &c., connected with the lamp inside the ionisation chamber might act as sources of impurities in Lenard and Ramsauer's sense.

If we consider the quantum theory of radiation to apply to ionisation of gases by light, then the energy available in the quantum, *hn*, must exceed the work V_0e required to separate an electron from a molecule. Palmer's experiments (*Phys. Rev.*, xxxii., p. 1, 1911) may perhaps be taken to indicate that the oxygen accounts for most of the ionisation in air. Taking the longest wave-length which ionises air to be λ1350, and $h=6{\cdot}55\times10^{-27}$, and $e=4{\cdot}65\times10^{-10}$, we get $V_0=9{\cdot}2$ volts. Now the ionising potential for oxygen according to Frank and Hertz is 9·0 volts. To maintain that λ1800 is nearer to the long wave-length limit implies that the quantum theory is not applicable to ionisation by light, for there is no reason to doubt the accuracy of the experiments of Frank and Hertz.

A. LL. HUGHES.

Cavendish Laboratory, Cambridge.

The Microtropometer.

MANY roads to progress in physical investigation are brought to an abrupt end through the lack of measuring instruments of sufficient sensitiveness. In the attempt to bridge over one of these disabling chasms the writer was led to the following device, which appears capable of some development. The principle can be illustrated with reference to a particular case. Suppose we have a Boys's radio-micrometer, which we will call the secondary instrument. If we project on to the vane of this an image of a Nernst lamp filament the beam of light from the mirror of the instrument may be deflected through one thousand scale divisions. Suppose now that the image of the filament is 1 mm. wide, and that it is projected by the mirror of another radio-micrometer, which we may call the primary one.

It is evident that a movement of this primary image through a distance of 1 mm. can produce a movement of the beam of light from the secondary instrument through 1000 mm. Hence, a movement of the image of the filament cast by the primary instrument through 0·001 mm. would give a deflection of about 1 mm., and a movement of the primary image through 1/1,000,000 mm. would move the secondary image through one-thousandth of a mm. If now the secondary instrument be made to throw a similar image on to a tertiary radio-micrometer, the motion will again be magnified one thousand times, so that an original movement of a millionth of a mm. produces a final movement of 1 mm. Evidently by increasing the number of instruments in arithmetical progression we increase the magnification in geometrical progression.

I have applied the method to two radio-micrometers with very satisfactory results. The principle, however, can be applied to any instrument in which a beam of light is used as an indicator—*e.g.* the primary instrument may be a galvanometer, an electrometer, a double-thread-suspension mirror, or a string-galvanometer (in the last case the image of the string taking the place of the image of the filament). The secondary instruments may be radio-micrometers, thermo-couples, bolometers, selenium cells, or other detectors of radiation. It will appear, therefore, that the principle is one capable of wide application to cases in which great sensitiveness of measurement is required—from wireless telegraphy to physiology. In fact, we may say that any existing instrument which uses light as an indicator can be made more sensitive.

Practical difficulties arise from the impossibility of obtaining any instrument with absolutely constant zero; moreover, fluctuations in the intensity of the energy stream from the source of radiation, represented by the Nernst filament, would cause trouble

©1913 Nature Publishing Group

where great magnifications were desired. This last difficulty might to some extent be surmounted by using as a secondary instrument the two strips of a bolometer, so that fluctuations would make proportional changes in both strips, whereas a deflection gives a differential effect. Experiments, however, on the application of the principle are in progress.

G. A. SHAKESPEAR.

The University, Birmingham, June 24.

Wireless Antennæ.

IN recent issues of NATURE several correspondents, in referring to the fact that a metal bedstead or a few wires stretched a few feet above the ground will make a wireless antenna, have overlooked a most important point, viz. that with such an antenna the ordinary methods of tuning are quite useless.

A piece of wire netting suspended a few feet above the ground makes a most effective aërial, and enables one to receive loud signals from long-distance stations, but signals from Eiffel Tower, Cleethorpes, &c., will all be mixed up, and the ordinary tuner will not separate them effectively. It seems to me that this proves that the usual theory of the waves travelling through space in the air above the earth's surface and being cut by the aërial does not *wholly* account for the facts.

Wireless signals that are feeble when the surface of the earth is dry, becoming much stronger after rain, and the well-known fact that these waves travel much better over sea than over land, all seem to indicate that the aërial waves are *at least* supplemented by waves that travel along the surface of the earth, and that the wire netting, bedsteads, &c., act as counter capacities, allowing these waves to flow from the earth through the receiver. The capacity of a small piece of wire netting near the ground is much greater than a very extensive aërial high up.

Canterbury. A. LANDER.

The Occurrence of "Anomalocera pattersoni," Temp., in Mounts Bay.

IT is stated by Mr. L. H. Gough ("Report of the Plankton of the English Channel," 1903), that the copepod *Anomalocera pattersoni*, R. Temp., may generally be regarded as an oceanic species. Gough's observations tended to show that Gran was correct in assuming this to be the case, although Cleve considered the species to be neritic. Sars, in his "Crustacea of Norway," speaks of the occurrence of this copepod "off the south and west coasts of Norway, generally congregated in great shoals," but throughout the reports published in connection with the international fishery investigations, no record occurs of its similar abundant occurrence in the English Channel.

It is interesting to record, therefore, that during an intensive survey of the planktological conditions of Mounts Bay, performed from the s.y. *Mera* on June 2, an extensive shoal of the species was met with at the surface on a position 50° 4′ 20″ N. × 5° 27′ 55″ W. The product of a five minutes' haul with the coarse tow-net amounted to 34·75 c.c. of material, of which at least three-fourths was made up of Anomalocera, both in its adult and developmental stages. The visitation of this copepod, in association with several other oceanic forms of phyto- and zooplankton, to a comparatively close inshore position may possibly serve as an indication of somewhat abnormal hydrographical conditions, but unfortunately no physical observations are yet to hand to enable a comparison to be formed.

HAROLD SWITHINBANK.
S.y. *Mera*, R.Y.S. G. E. BULLEN.

Artificial Hiss.

LORD RAYLEIGH'S "sound more like an *f* than an *s*" (NATURE, May 29, p. 319) is due, according to Köhler's observations (*Zeits. f. Psych.*, 64, p. 93), to a slightly too high pitch. A Galton whistle, set for a tone of 8400 v.d., will give a pure *s*.

E. B. TITCHENER.

The Psychological Laboratory, Cornell University, Ithaca, N.Y.

THE BICENTENARY OF THE BOTANIC GARDEN OF ST. PETERSBURG.

THE bicentenary of the foundation of the Imperial Botanic Garden at St. Petersburg was celebrated with unusual pomp on June 24. The history of the garden, its share in the botanical exploration of north, central, and east Asia, and the practical completion of its reorganisation on modern and truly liberal lines justified the festive character of the proceedings. They were initiated on the eve of the jubilee by a special service, attended by the staff of the garden, in the Peter-Paul's Cathedral, and the laying down of a wreath of roses at the tomb of Peter the Great, the founder of the garden, and on the following morning by a little shower of honours for the director and his staff.

The principal ceremony took place in the afternoon in a building attached to the old herbarium, in the presence of a brilliant gathering, including the Princes Peter and Alexander of Oldenburg, Prince Gagarin, representing Princess Eugenia of Oldenburg, under whose patronage the garden is placed, the Ministers of Agriculture, Commerce, Justice, Public Cults, and Marine, and other prominent officials, and a large number of delegates from the Russian academies, universities, botanic gardens, and kindred institutions, and from foreign countries. The solemn meeting was preceded by a Te Deum, and formally opened by the Minister of Agriculture, Mr. Kriwoshein. A rescript from the Emperor was read, in which it was announced that the garden would henceforth be known as the Imperial Botanic Garden of Peter the Great. Then followed an eloquent address by the director of the garden, Prof. A. Fischer von Waldheim, in which he traced the history of the garden and its present organisation and object; the reception of the delegates, and the presentation of medals and souvenirs commemorating the jubilee.

The next day was reserved for the inspection of the garden, museum, and laboratories, and in the afternoon a visit to Peterhof, where the guests experienced an exceptionally cordial reception. In the evening the festivities came to a conclusion with a gala performance in the opera house in the Narodni Dom.

Most of the botanical gardens and many of the other botanical institutions sent their congratulations, whilst Bulgaria, Denmark, England (Kew and Chelsea), Germany, and Sweden were represented by delegates.

The garden was founded by Peter the Great about 1713, so that it is practically as old as the

©1913 Nature Publishing Group